

A STIELTJES FUNCTION INVOLVING THE LOGARITHMIC FUNCTION AND AN APPLICATION

FENG QI AND XIAO-JING ZHANG

ABSTRACT. In the paper, the authors establish an integral representation of the holomorphic extension to the cut plane of a function involving the logarithmic function, and prove that the function involving the logarithmic function is a Stieltjes function. By applying this integral representation, the authors discover an integral representation for Bernoulli numbers of the second kind and obtain the complete monotonicity of the sequence of Bernoulli numbers of the second kind.

1. INTRODUCTION

Let us recall some definitions we will use in this paper.

Definition 1.1 ([9, Chapter XIII] and [21, Chapter IV]). A function f is said to be completely monotonic on an interval I if f has derivatives of all orders on I and

$$(-1)^n f^{(n)}(x) \geq 0 \quad (1)$$

for $x \in I$ and $n \geq 0$.

Definition 1.2 ([19, p. 19, Definition 2.1]). A Stieltjes function is a function $f : (0, \infty) \rightarrow [0, \infty)$ which can be written in the form

$$f(x) = \frac{a}{x} + b + \int_0^\infty \frac{1}{s+x} d\mu(s), \quad (2)$$

where $a, b \geq 0$ are nonnegative constants and μ is a nonnegative measure on $(0, \infty)$ such that

$$\int_0^\infty \frac{1}{1+s} d\mu(s) < \infty.$$

Definition 1.3 ([2, p. 457]). A function $f : I \subseteq (0, \infty) \rightarrow [0, \infty)$ is called a Bernstein function on I if $f(x)$ has derivatives of all orders and $f'(x)$ is completely monotonic on I .

Definition 1.4 ([3, 12, 13, 16]). A function $f(x)$ is said to be logarithmically completely monotonic on an interval $I \subseteq \mathbb{R}$ if it has derivatives of all orders on I and its logarithm $\ln f(x)$ satisfies

$$(-1)^k [\ln f(x)]^{(k)} \geq 0 \quad (3)$$

for $k \in \mathbb{N}$ on I .

2010 *Mathematics Subject Classification.* Primary 30E20; Secondary 11B68, 11B83, 26A48, 44A15, 47B06.

Key words and phrases. Integral representation; Stieltjes function; Cauchy integral formula; Logarithmic function; Bernoulli numbers of the second kind; Completely monotonic sequence.

In what follows, we use $\mathcal{C}[(0, \infty)]$, $\mathcal{L}[(0, \infty)]$, $\mathcal{B}[(0, \infty)]$, and $\mathcal{S}$ to denote the classes of completely monotonic functions on $(0, \infty)$, of logarithmically completely monotonic functions on $(0, \infty)$, of Bernstein functions on $(0, \infty)$, and of Stieltjes functions respectively.

These classes of functions have the following properties and relations.

Proposition 1.1 ([21, p. 161, Theorem 12b]). *A necessary and sufficient condition that $f(x)$ should be completely monotonic for $0 < x < \infty$ is that*

$$f(x) = \int_0^\infty e^{-xt} d\alpha(t), \quad (4)$$

where $\alpha(t)$ is non-decreasing and the integral converges for $0 < x < \infty$.

Proposition 1.2 ([19, p. 15, Theorem 3.2]). *A function $f : (0, \infty) \rightarrow \mathbb{R}$ is a Bernstein function if and only if it admits the representation*

$$f(x) = a + bx + \int_0^\infty (1 - e^{-xt}) d\mu(t), \quad (5)$$

where $a, b \geq 0$ and μ is a measure on $(0, \infty)$ satisfying

$$\int_0^\infty \min\{1, t\} d\mu(t) < \infty.$$

Proposition 1.3 ([4, 8, 12, 13]).

$$\mathcal{S} \setminus \{0\} \subset \mathcal{L}[(0, \infty)] \subset \mathcal{C}[(0, \infty)]. \quad (6)$$

Proposition 1.4 ([6, pp. 161–162, Theorem 3] and [19, p. 45, Proposition 5.17]). *If $f(x)$ is a Bernstein function on an interval I , then $\frac{1}{f(x)}$ is logarithmically completely monotonic on I .*

Proposition 1.5 ([19, p. 19, Theorem 3.6]). *Let f be a positive function on $(0, \infty)$. Then the following assertions are equivalent.*

- (1) *The function f is a Bernstein function.*
- (2) *For every completely monotonic function g , the composite $g \circ f$ is also completely monotonic.*
- (3) *For every $u > 0$, the function e^{-uf} is completely monotonic.*

For more information on these four classes of functions, please refer to [8, 9, 10, 11, 17, 18, 19, 21] and plenty of references cited therein.

By the way, the preprints [13, 16] were extended, revised, and formally published in [12, 14, 15] respectively.

As usual, we use $\ln x$ for the logarithmic function having base e and applied to the positive argument $x > 0$. Further, it is general knowledge that the principal branch of the holomorphic extension of $\ln x$ from the open half-line $(0, \infty)$ to the cut plane $\mathcal{A} = \mathbb{C} \setminus (-\infty, 0]$ is denoted by

$$\ln z = \ln |z| + i \arg z,$$

where $-\pi < \arg z < \pi$ and $i = \sqrt{-1}$. See related texts in [7].

By a standard argument, we can reveal that the function $\ln(1+x)$ is a Bernstein function on $(0, \infty)$. Therefore, by Proposition 1.4, it is derived that the function

$\frac{1}{\ln(1+x)}$ is logarithmically completely monotonic on $(0, \infty)$. This conclusion may also be directly verified as follows. In [1, p. 230, 5.1.32], it is listed that

$$\ln \frac{b}{a} = \int_0^\infty \frac{e^{-au} - e^{-bu}}{u} du. \quad (7)$$

As a result, we have

$$\ln[\ln(1+x)] = \int_0^\infty \frac{e^{-u} - e^{-u \ln(1+x)}}{u} du = \int_0^\infty \frac{e^{-u} - (1+x)^{-u}}{u} du \quad (8)$$

and

$$\frac{1}{(1+x) \ln(1+x)} = \int_0^\infty \frac{1}{(1+x)^{u+1}} du, \quad x > 0. \quad (9)$$

Thus, the function $\frac{1}{(1+x) \ln(1+x)}$ is completely monotonic on $(0, \infty)$. Equivalently and strongly speaking, the function $\frac{1}{\ln(1+x)}$ is logarithmically completely monotonic on $(0, \infty)$.

More strongly, it is known that the function $\frac{1}{\ln(1+x)}$ is a Stieltjes function and has the integral representation

$$\frac{1}{\ln(1+z)} = \frac{1}{z} + \int_1^\infty \frac{1}{[\ln(t-1)]^2 + \pi^2} \frac{dt}{z+t}, \quad z \in \mathcal{A}, \quad (10)$$

which is a corrected version of the equation (34) in [5, p. 2130].

We are now interested in considering

$$F(x) = \frac{1}{\ln(1+x)} - \frac{1}{(1+x) \ln(1+x)} = \frac{x}{(1+x) \ln(1+x)} \quad (11)$$

on $(0, \infty)$, which is, by (6), a difference of two (logarithmically) completely monotonic functions. Here we remark that a sum or product of any finitely many completely monotonic functions is still completely monotonic and that a product of any finitely many logarithmically completely monotonic functions is also logarithmically completely monotonic, see [14], but a difference or a quotient of any two (logarithmically) completely monotonic functions is not always a (logarithmically) completely monotonic function. So, to consider the function $F(x)$ is meaningful and significant.

It is easy to see that the principal branch of the holomorphic extension of $F(x)$ to the cut plane $\mathcal{A}$ is

$$F(z) = \frac{z}{(1+z) \ln(1+z)}, \quad z \in \mathcal{A}. \quad (12)$$

The first goal of this paper is to establish an integral representation of $F(z)$ and, as a consequence, to reveal that $F(x)$ is a Stieltjes function.

The first main result may be stated as the following theorem.

Theorem 1.1. *The function $F(z)$ defined by (12) has the integral representation*

$$F(z) = \int_1^\infty \frac{t}{(t-1)\{[\ln(t-1)]^2 + \pi^2\}} \frac{dt}{z+t}, \quad z \in \mathcal{A}. \quad (13)$$

Consequently, the function $F(x)$ is a Stieltjes function.

Bernoulli numbers of the second kind $b_0, b_1, b_2, \dots, b_n, \dots$ are defined by

$$\frac{x}{\ln(1+x)} = \sum_{n=0}^\infty b_n x^n. \quad (14)$$

They are also known as Cauchy numbers, Gregory coefficients, or logarithmic numbers. See [20]. The first few Bernoulli numbers of the second kind are

$$b_0 = 1, \quad b_1 = \frac{1}{2}, \quad b_2 = -\frac{1}{12}, \quad b_3 = \frac{1}{24}, \quad b_4 = -\frac{19}{720}, \quad b_5 = \frac{3}{160}. \quad (15)$$

The second goal of this paper is to apply the integral representation (13) to discover an integral representation (16) for Bernoulli numbers of the second kind.

Theorem 1.2. *Bernoulli numbers of the second kind b_n may be calculated by*

$$b_n = (-1)^{n+1} \int_1^\infty \frac{1}{\{\ln(t-1)\}^2 + \pi^2} t^n dt, \quad n \in \mathbb{N}. \quad (16)$$

We recall from [21, p. 108, Definition 4] that a sequence $\{\mu_n\}_0^\infty$ is said to be completely monotonic if its elements are non-negative and its successive differences are alternatively non-negative, that is

$$(-1)^k \Delta^k \mu_n \geq 0 \quad (17)$$

for $n, k \geq 0$, where

$$\Delta^k \mu_n = \sum_{m=0}^k (-1)^m \binom{k}{m} \mu_{n+k-m}. \quad (18)$$

Theorem 4a in [21, p. 108] reads that a necessary and sufficient condition that the sequence $\{\mu_n\}_0^\infty$ should have the expression

$$\mu_n = \int_0^1 t^n d\alpha(t) \quad (19)$$

for $n \geq 0$, where $\alpha(t)$ is non-decreasing and bounded for $0 \leq t \leq 1$, is that it should be completely monotonic.

The third goal of this paper is to, making use of the integral representation (16), obtain the complete monotonicity of the sequence $\{(-1)^n b_{n+1}\}_{n=0}^\infty$ of Bernoulli numbers of the second kind.

Theorem 1.3. *The sequence $\{(-1)^n b_{n+1}\}_{n=0}^\infty$ of Bernoulli numbers of the second kind is a completely monotonic sequence.*

2. LEMMAS

In order to prove Theorem 1.1, some lemmas are needed.

Lemma 2.1 (Cauchy integral formula [7, p. 113]). *Let D be a bounded domain with piecewise smooth boundary. If $f(z)$ is analytic on D , and $f(z)$ extends smoothly to the boundary of D , then*

$$f(z) = \frac{1}{2\pi i} \oint_{\partial D} \frac{f(w)}{w-z} dw, \quad z \in D. \quad (20)$$

Lemma 2.2. *For $z = \varepsilon e^{i\theta}$ with $\theta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ and $\varepsilon \in (0, 1)$, we have*

$$\lim_{\varepsilon \rightarrow 0^+} [zF(z)] = 0. \quad (21)$$

Proof. It is enough to show $|zF(z)| \rightarrow 0$ as $\varepsilon \rightarrow 0^+$.

For $z = \varepsilon e^{i\theta}$ and $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, we have

$$|zF(z)| = \left| \frac{(\varepsilon e^{i\theta})^2}{(1 + \varepsilon e^{i\theta}) \ln(1 + \varepsilon e^{i\theta})} \right|$$

$$\begin{aligned}
&= \frac{|\varepsilon^2 e^{2i\theta}|}{|1 + \varepsilon e^{i\theta}| |\ln |1 + \varepsilon e^{i\theta}| + i \arg(1 + \varepsilon e^{i\theta})|} \\
&\leq \frac{\varepsilon^2}{|1 + \varepsilon e^{i\theta}| |\ln |1 + \varepsilon e^{i\theta}||} \\
&= \frac{\varepsilon^2}{\sqrt{1 + \varepsilon^2 + 2\varepsilon \cos \theta} |\ln \sqrt{1 + \varepsilon^2 + 2\varepsilon \cos \theta}|}.
\end{aligned}$$

It is easy to see that

$$\begin{aligned}
&\lim_{\varepsilon \rightarrow 0^+} \frac{\varepsilon^2}{\sqrt{1 + \varepsilon^2 + 2\varepsilon \cos \theta} |\ln \sqrt{1 + \varepsilon^2 + 2\varepsilon \cos \theta}|} \\
&= 2 \lim_{\varepsilon \rightarrow 0^+} \frac{\varepsilon^2}{|\ln(1 + \varepsilon^2 + 2\varepsilon \cos \theta)|} \\
&= 2 \lim_{\varepsilon \rightarrow 0^+} \frac{\varepsilon^2}{|\varepsilon^2 + 2\varepsilon \cos \theta|} \\
&= 0.
\end{aligned}$$

Hence, it follows that $|zF(z)| \rightarrow 0$ as $\varepsilon \rightarrow 0^+$.

For $z = \pm i\varepsilon$, we have

$$\begin{aligned}
|zF(z)| &= \left| \frac{(\pm i\varepsilon)^2}{(1 \pm i\varepsilon) \ln(1 \pm i\varepsilon)} \right| \\
&= \frac{\varepsilon^2}{|1 \pm i\varepsilon| |\ln |1 \pm i\varepsilon| + i \arg(1 \pm i\varepsilon)|} \\
&= \frac{\varepsilon^2}{\sqrt{1 + \varepsilon^2} \sqrt{(\ln |1 \pm i\varepsilon|)^2 + [\arg(1 \pm i\varepsilon)]^2}} \\
&= \frac{\varepsilon^2}{\sqrt{1 + \varepsilon^2} \sqrt{[\ln(1 + \varepsilon^2)]^2/4 + (\arctan \varepsilon)^2}} \\
&= \frac{\varepsilon}{\sqrt{1 + \varepsilon^2} \sqrt{\{[\ln(1 + \varepsilon^2)]/\varepsilon\}^2/4 + [(\arctan \varepsilon)/\varepsilon]^2}} \\
&\rightarrow \frac{0}{\sqrt{1 + 0^2} \sqrt{0^2/4 + 1^2}} \\
&= 0.
\end{aligned}$$

The proof is complete. $\square$

Lemma 2.3. *As $r \rightarrow \infty$, the function $F(z)$ tends to 0 uniformly for $\theta \in (-\pi, \pi)$ and $z = re^{i\theta}$.*

Proof. It is not difficult to see that

$$\begin{aligned}
|F(z)| &= \left| \frac{re^{i\theta}}{(1 + re^{i\theta}) \ln(1 + re^{i\theta})} \right| \\
&= \frac{r}{|1 + re^{i\theta}| |\ln |1 + re^{i\theta}| + i \arg(1 + re^{i\theta})|} \\
&\leq \frac{r}{|1 + re^{i\theta}| |\ln |1 + re^{i\theta}||} \\
&= \frac{r}{\sqrt{1 + r^2 + 2r \cos \theta} |\ln \sqrt{1 + r^2 + 2r \cos \theta}|}
\end{aligned}$$

$$\begin{aligned} &\leq \frac{r}{(r-1)\ln(r-1)} \\ &\rightarrow 0 \end{aligned}$$

as $r \rightarrow \infty$. The proof is thus complete. $\square$

Lemma 2.4. *For $t \in (0, \infty)$, we have*

$$\lim_{\varepsilon \rightarrow 0^+} \Im F(-t + i\varepsilon) = \begin{cases} 0, & 0 < t < 1; \\ -\frac{\pi t}{(t-1)\{\ln(t-1)^2 + \pi^2\}}, & t > 1; \\ \infty, & t = 1. \end{cases} \quad (22)$$

Proof. For $t \in (0, \infty)$ and $\varepsilon \in (0, 1)$, we have


$$\begin{aligned} F(-t + i\varepsilon) &= \frac{-t + i\varepsilon}{(1-t+i\varepsilon)\ln(1-t+i\varepsilon)} \\ &= \frac{-t + i\varepsilon}{(1-t+i\varepsilon)[\ln|1-t+i\varepsilon| + i\arg(1-t+i\varepsilon)]} \\ &= \begin{cases} \frac{-t + i\varepsilon}{(1-t+i\varepsilon)[\ln|1-t+i\varepsilon| + i\arctan\frac{\varepsilon}{1-t}]}, & 1-t > 0 \\ \frac{-t + i\varepsilon}{(1-t+i\varepsilon)[\ln|1-t+i\varepsilon| + i(\pi + \arctan\frac{\varepsilon}{1-t})]}, & 1-t < 0 \\ \frac{-1 + i\varepsilon}{i\varepsilon(\ln\varepsilon + i\frac{\pi}{2})}, & 1-t = 0 \end{cases} \\ &\rightarrow \begin{cases} \frac{-t}{(1-t)\ln|1-t|}, & 1-t > 0 \\ \frac{-t}{(1-t)[\ln|1-t| + \pi i]}, & 1-t < 0 \\ \frac{-1}{i\lim_{\varepsilon \rightarrow 0^+} [\varepsilon(\ln\varepsilon + i\frac{\pi}{2})]}, & 1-t = 0 \end{cases} \\ &= \begin{cases} \frac{t}{(t-1)\ln(1-t)}, & 0 < t < 1 \\ \frac{t[\ln(t-1) - \pi i]}{(t-1)\{\ln(t-1)^2 + \pi^2\}}, & t > 1 \\ \infty, & t = 1 \end{cases} \end{aligned}$$

as $\varepsilon \rightarrow 0^+$. The proof is thus complete. $\square$

3. PROOFS OF THEOREMS 1.1, 1.2, AND 1.3

Now we are in a position to prove Theorems 1.1, 1.2, and 1.3 in sequence.

Proof of Theorem 1.1. For any fixed point $z \in \mathcal{A}$, choose $0 < \varepsilon < 1$ and $r > 0$ such that $0 < \varepsilon < |z| < r$, and consider the positively oriented contour $C(\varepsilon, r)$ in $\mathcal{A}$ consisting of the half circle $z = \varepsilon e^{i\theta}$ for $\theta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ and the half lines $z = x \pm i\varepsilon$ for $x \leq 0$ until they cut the circle $|z| = r$, which close the contour at the points $-r(\varepsilon) \pm i\varepsilon$, where $0 < r(\varepsilon) \rightarrow r$ as $\varepsilon \rightarrow 0$. See Figure 1. By Lemma 2.1 we have

FIGURE 1. The contour $C(\varepsilon, r)$

$$\begin{aligned}
 F(z) &= \frac{1}{2\pi i} \oint_{C(\varepsilon, r)} \frac{F(w)}{w - z} dw \\
 &= \frac{1}{2\pi i} \left[\int_{\pi/2}^{-\pi/2} \frac{i\varepsilon e^{i\theta} F(\varepsilon e^{i\theta})}{\varepsilon e^{i\theta} - z} d\theta + \int_{-r(\varepsilon)}^0 \frac{F(x + i\varepsilon)}{x + i\varepsilon - z} dx \right. \\
 &\quad \left. + \int_0^{-r(\varepsilon)} \frac{F(x - i\varepsilon)}{x - i\varepsilon - z} dx + \int_{\arg[-r(\varepsilon) - i\varepsilon]}^{\arg[-r(\varepsilon) + i\varepsilon]} \frac{ire^{i\theta} F(re^{i\theta})}{re^{i\theta} - z} d\theta \right].
 \end{aligned} \tag{23}$$

By Lemma 2.2, it follows that

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\pi/2}^{-\pi/2} \frac{i\varepsilon e^{i\theta} F(\varepsilon e^{i\theta})}{\varepsilon e^{i\theta} - z} d\theta = 0. \tag{24}$$

In virtue of Lemma 2.3, it can be derived that

$$\lim_{\substack{\varepsilon \rightarrow 0^+ \\ r \rightarrow \infty}} \int_{\arg[-r(\varepsilon) - i\varepsilon]}^{\arg[-r(\varepsilon) + i\varepsilon]} \frac{ire^{i\theta} F(re^{i\theta})}{re^{i\theta} - z} d\theta = \lim_{r \rightarrow \infty} \int_{-\pi}^{\pi} \frac{ire^{i\theta} F(re^{i\theta})}{re^{i\theta} - z} d\theta = 0. \tag{25}$$

Making use of the obvious fact that $F(\bar{z}) = \overline{F(z)}$ and Lemma 2.4 yields that

$$\begin{aligned}
 &\int_{-r(\varepsilon)}^0 \frac{F(x + i\varepsilon)}{x + i\varepsilon - z} dx + \int_0^{-r(\varepsilon)} \frac{F(x - i\varepsilon)}{x - i\varepsilon - z} dx \\
 &= \int_{-r(\varepsilon)}^0 \left[\frac{F(x + i\varepsilon)}{x + i\varepsilon - z} - \frac{F(x - i\varepsilon)}{x - i\varepsilon - z} \right] dx
 \end{aligned}$$

$$\begin{aligned}
&= \int_{-r(\varepsilon)}^0 \frac{(x - i\varepsilon - z)F(x + i\varepsilon) - (x + i\varepsilon - z)F(x - i\varepsilon)}{(x + i\varepsilon - z)(x - i\varepsilon - z)} dx \\
&= \int_{-r(\varepsilon)}^0 \frac{(x - z)[F(x + i\varepsilon) - F(x - i\varepsilon)] - i\varepsilon[F(x - i\varepsilon) + F(x + i\varepsilon)]}{(x + i\varepsilon - z)(x - i\varepsilon - z)} dx \\
&= 2i \int_{-r(\varepsilon)}^0 \frac{(x - z)\Im F(x + i\varepsilon) - \varepsilon\Re F(x + i\varepsilon)}{(x + i\varepsilon - z)(x - i\varepsilon - z)} dx \\
&\rightarrow 2i \int_{-r}^0 \frac{\lim_{\varepsilon \rightarrow 0^+} \Im F(x + i\varepsilon)}{x - z} dx \\
&= -2i \int_0^r \frac{\lim_{\varepsilon \rightarrow 0^+} \Im F(-t + i\varepsilon)}{t + z} dt \\
&= -2i \int_0^\infty \frac{\lim_{\varepsilon \rightarrow 0^+} \Im F(-t + i\varepsilon)}{t + z} dt \\
&= 2\pi i \int_1^\infty \frac{t}{(t - 1)\{\ln(t - 1)^2 + \pi^2\}} \frac{dt}{t + z} \tag{26}
\end{aligned}$$

as $\varepsilon \rightarrow 0^+$ and $r \rightarrow \infty$. Substituting equations (24), (25), and (26) into (23) and simplifying produce the integral representation (13). The proof of Theorem 1.1 is complete. $\square$

Proof of Theorem 1.2. By the integral representation (13), we have

$$\frac{x}{\ln(1+x)} = \int_1^\infty \frac{t}{(t-1)\{\ln(t-1)^2 + \pi^2\}} \frac{1+x}{x+t} dt \tag{27}$$

and

$$\begin{aligned}
\left[\frac{x}{\ln(1+x)} \right]^{(k)} &= \int_1^\infty \frac{t}{(t-1)\{\ln(t-1)^2 + \pi^2\}} \left(\frac{1+x}{x+t} \right)^{(k)} dt \\
&= \int_1^\infty \frac{t}{(t-1)\{\ln(t-1)^2 + \pi^2\}} \left(1 + \frac{1-t}{x+t} \right)^{(k)} dt \\
&= (-1)^{k+1} k! \int_1^\infty \frac{t}{[\ln(t-1)]^2 + \pi^2} \frac{1}{(x+t)^{k+1}} dt
\end{aligned} \tag{28}$$

for $k \in \mathbb{N}$. On the other hand, by (14), we also have

$$\left[\frac{x}{\ln(1+x)} \right]^{(k)} = \sum_{n=k}^\infty b_n \frac{n!}{(n-k)!} x^{n-k}. \tag{29}$$

Combining (28) with (29) leads to

$$\sum_{n=k}^\infty b_n \frac{n!}{(n-k)!} x^{n-k} = (-1)^{k+1} k! \int_1^\infty \frac{t}{[\ln(t-1)]^2 + \pi^2} \frac{1}{(x+t)^{k+1}} dt. \tag{30}$$

Letting $x \rightarrow 0^+$ on both sides of (30) yields the formula (16). The proof of Theorem 1.2 is completed. $\square$

Proof of Theorem 1.3. From (16), it follows that for $n \in \mathbb{N}$

$$(-1)^{n+1} b_n = \int_1^\infty \frac{1}{\{\ln(t-1)\}^2 + \pi^2} t^n dt$$

$$\begin{aligned}
&= \int_1^0 \frac{1}{\{[\ln(1/s-1)]^2 + \pi^2\}} s^n d\left(\frac{1}{s}\right) \\
&= \int_0^1 \frac{1}{\{[\ln(1/s-1)]^2 + \pi^2\}} s^{n-2} ds \\
&= \int_0^1 \frac{1}{s\{[\ln(1/s-1)]^2 + \pi^2\}} s^{n-1} ds \\
&\triangleq c_{n-1}.
\end{aligned}$$

Since $c_0 = b_1 = \frac{1}{2}$ and the function $\frac{1}{s\{[\ln(1/s-1)]^2 + \pi^2\}}$ is positive on $(0, 1)$, then the complete monotonicity of the sequence $\{c_n\}_0^\infty$ is readily obtained. The required result is thus proved. The proof of Theorem 1.3 is complete. $\square$

REFERENCES

- [1] M. Abramowitz and I. A. Stegun (Eds), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series **55**, 9th printing, Washington, 1970.
- [2] H. Alzer and C. Berg, *Some classes of completely monotonic functions*, Ann. Acad. Sci. Fenn. Math. **27** (2002), no. 2, 445–460.
- [3] R. D. Atanassov and U. V. Tsoukrovski, *Some properties of a class of logarithmically completely monotonic functions*, C. R. Acad. Bulgare Sci. **41** (1988), no. 2, 21–23.
- [4] C. Berg, *Integral representation of some functions related to the gamma function*, Mediterr. J. Math. **1** (2004), no. 4, 433–439; Available online at <http://arxiv.org/abs/math/0411550>.
- [5] C. Berg and H. L. Pedersen, *A one-parameter family of Pick functions defined by the Gamma function and related to the volume of the unit ball in n-space*, Proc. Amer. Math. Soc. **139** (2011), no. 6, 2121–2132; Available online at <http://dx.doi.org/10.1090/S0002-9939-2010-10636-6>.
- [6] C.-P. Chen, F. Qi, and H. M. Srivastava, *Some properties of functions related to the gamma and psi functions*, Integral Transforms Spec. Funct. **21** (2010), no. 2, 153–164; Available online at <http://dx.doi.org/10.1080/10652460903064216>.
- [7] T. W. Gamelin, *Complex Analysis*, Undergraduate Texts in Mathematics, Springer, New York-Berlin-Heidelberg, 2001.
- [8] B.-N. Guo and F. Qi, *A property of logarithmically absolutely monotonic functions and the logarithmically complete monotonicity of a power-exponential function*, Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys. **72** (2010), no. 2, 21–30.
- [9] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht-Boston-London, 1993.
- [10] F. Qi, *Bounds for the ratio of two gamma functions*, J. Inequal. Appl. **2010** (2010), Article ID 493058, 84 pages; Available online at <http://dx.doi.org/10.1155/2010/493058>.
- [11] F. Qi, *Certain logarithmically N-alternating monotonic functions involving gamma and q-gamma functions*, Nonlinear Funct. Anal. Appl. **12** (2007), no. 4, 675–685.
- [12] F. Qi and C.-P. Chen, *A complete monotonicity property of the gamma function*, J. Math. Anal. Appl. **296** (2004), no. 2, 603–607; Available online at <http://dx.doi.org/10.1016/j.jmaa.2004.04.026>.
- [13] F. Qi and B.-N. Guo, *Complete monotonicities of functions involving the gamma and digamma functions*, RGMIA Res. Rep. Coll. **7** (2004), no. 1, Art. 8, 63–72; Available online at <http://rgmia.org/v7n1.php>.
- [14] F. Qi and B.-N. Guo, *Some logarithmically completely monotonic functions related to the gamma function*, J. Korean Math. Soc. **47** (2010), no. 6, 1283–1297; Available online at <http://dx.doi.org/10.4134/JKMS.2010.47.6.1283>.
- [15] F. Qi, B.-N. Guo, and C.-P. Chen, *Some completely monotonic functions involving the gamma and polygamma functions*, J. Aust. Math. Soc. **80** (2006), 81–88; Available online at <http://dx.doi.org/10.1017/S1446788700011393>.

- [16] F. Qi, B.-N. Guo, and C.-P. Chen, *Some completely monotonic functions involving the gamma and polygamma functions*, RGMIA Res. Rep. Coll. **7** (2004), no. 1, Art. 5, 31–36; Available online at <http://rgmia.org/v7n1.php>.
- [17] F. Qi and Q.-M. Luo, *Bounds for the ratio of two gamma functions—From Wendel's and related inequalities to logarithmically completely monotonic functions*, Banach J. Math. Anal. **6** (2012), no. 2, 132–158.
- [18] F. Qi, C.-F. Wei, and B.-N. Guo, *Complete monotonicity of a function involving the ratio of gamma functions and applications*, Banach J. Math. Anal. **6** (2012), no. 1, 35–44.
- [19] R. L. Schilling, R. Song, and Z. Vondraček, *Bernstein Functions*, de Gruyter Studies in Mathematics **37**, De Gruyter, Berlin, Germany, 2010.
- [20] E. W. Weisstein, *Bernoulli Number of the Second Kind*, From MathWorld—A Wolfram Web Resource, Available online at <http://mathworld.wolfram.com/BernoulliNumberoftheSecondKind.html>.
- [21] D. V. Widder, *The Laplace Transform*, Princeton University Press, Princeton, 1946.

(Qi) DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN CITY, 300387, CHINA

E-mail address: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com

URL: <http://qifeng618.wordpress.com>

(Zhang) DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN CITY, 300387, CHINA

E-mail address: xiao.jing.zhang@qq.com